

## TOPIC 3: Solving Equations for One Variable

The procedure we use to solve radical equations can also be used to solve a given equation for a specified variable. We illustrate the process with one last example.

### Example 4: Escape Speed

The speed required for an object to escape from the gravitational pull of a planet is called the **escape speed** of the planet. The escape speed is given by the equation

$v_e = \sqrt{\frac{2GM}{r}}$ , where  $v_e$  is the escape speed,  $G$  is the universal gravitation constant,  $M$  is the mass of the planet, and  $r$  is the radius of the planet. Solve this equation for  $r$ .

#### Solution

We follow the same procedure for solving radical equations.

$$v_e = \sqrt{\frac{2GM}{r}} \quad \text{The radical expression is already isolated.}$$

$$v_e^2 = \frac{2GM}{r} \quad \text{Square both sides to eliminate the radical.}$$

$$r = \frac{2GM}{v_e^2} \quad \text{Solve for } r.$$

## 2.6 EXERCISES

### PRACTICE

Solve the following radical equations. See Example 2.

1.  $\sqrt{4-x} - x = 2$

2.  $\sqrt{3y+4} + \sqrt{5y+6} = 2$

3.  $\sqrt{3-3x} - 3 = \sqrt{3x+2}$

4.  $\sqrt{x^2 - 4x + 5} - x + 2 = 0$

5.  $\sqrt{x^2 - 4x + 4} + 2 = 3x$

6.  $\sqrt{50+7s} - s = 8$

7.  $\sqrt[3]{3-2x} - \sqrt[3]{x+1} = 0$

8.  $\sqrt[4]{x^2 - x} = \sqrt[4]{x-1}$

9.  $\sqrt[4]{2x+3} = -1$

10.  $\sqrt{11x+3} + 4x = 18$

11.  $\sqrt{2b-1} + 3 = \sqrt{10b-6}$

12.  $\sqrt{5x+5} = \sqrt{4x-7} + 2$

13.  $\sqrt{x+10} + 1 = x-1$

14.  $\sqrt{x+1} + 10 = x-1$

15.  $\sqrt{x^2 - 10} - 1 = x+1$

16.  $\sqrt[3]{5x^2 - 14x} = -2$

17.  $\sqrt[5]{7t^2 + 2t} = \sqrt[5]{5t^2 + 4}$

18.  $\sqrt[3]{y^3 - 7y + 2} = \sqrt[3]{2-3y}$

19.  $\sqrt{14y^2 - 18y + 4} + 2 = 2y$

20.  $\sqrt{9x+4} = \sqrt{7x+1} + 1$

21.  $\sqrt{4z+41}+3=z+2$

Solve the following equations. See Example 3.

22.  $(x+3)^{\frac{1}{4}}+2=0$

23.  $(2x-5)^{\frac{1}{4}}=(x-1)^{\frac{1}{4}}$

24.  $(2x-1)^{\frac{2}{3}}=x^{\frac{1}{3}}$

25.  $(3y^2+9y-5)^{\frac{1}{2}}=y+3$

26.  $(3x-5)^{\frac{1}{5}}=(x+1)^{\frac{1}{5}}$

27.  $w^{\frac{3}{5}}+8=0$

28.  $z^{\frac{4}{3}}-\frac{16}{81}=0$

29.  $x^{\frac{2}{3}}-\frac{25}{49}=0$

30.  $(x^2+21)^{\frac{-3}{2}}=\frac{1}{125}$

31.  $(x-2)^{\frac{2}{3}}=(14-x)^{\frac{1}{3}}$

32.  $(x^2+7)^{\frac{-3}{2}}=\frac{1}{64}$

33.  $(y-2)^{\frac{2}{3}}=(13y-66)^{\frac{1}{3}}$

Solve the following formulas for the indicated variable. See Example 4.

34. The formula  $T=2\pi\sqrt{\frac{l}{g}}$  gives the period  $T$  of a pendulum of length  $l$ . Solve this formula for  $l$ .

35. The formula  $c=\sqrt{a^2+b^2}$  gives the length of the hypotenuse  $c$  of a right triangle. Solve this formula for  $a$ .

36. Einstein's Theory of Relativity states that  $E=mc^2$ . Solve this equation for  $c$ .

37. The formula  $\omega=\sqrt{\frac{k}{m}}$  gives the angular frequency  $\omega$  of a mass  $m$  suspended from a spring of spring constant  $k$ . Solve this formula for  $m$ .

38. The formula  $V=\frac{4}{3}\pi r^3$  gives the volume of a sphere with radius  $r$ . Solve the equation for  $r$ .

39. The formula  $F=\frac{mv^2}{r}$  gives the force on an object in circular motion. Solve the equation for  $v$ .

40. The formula for lateral acceleration, used in automobiles, is  $a=\frac{1.227r}{t^2}$ . Solve this equation for  $t$ .

41. According to one guideline regarding body mass index, a healthy mass for an adult male can be found using the formula  $m=23h^2$ , where  $m$  is expressed in kilograms and  $h$  in meters. Solve this equation for  $h$ .

42. Kepler's Third Law is  $T^2=\frac{4\pi^2r^3}{GM}$ . It relates the period  $T$  of a planet to the radius  $r$  of its orbit and the sun's mass  $M$ . Solve this formula for  $r$ .

43. The equation  $r = \frac{2gm}{c^2}$  is the Schwarzschild Radius Formula used to find the radius of a black hole in space. Solve the equation for  $c$ .
44. The total mechanical energy of an object with mass  $m$  at height  $h$  in a closed system can be written as  $ME = \frac{1}{2}mv^2 + mgh$ . Solve for  $v$ , the velocity of the object, in terms of the given quantities.
45. Recall, the formula for the Pythagorean Theorem states that  $a^2 + b^2 = c^2$ . Solve this formula for  $b$ .
46. In a circuit with an AC power source, the total impedance  $Z$  depends on the resistance  $R$ , the capacitance  $C$ , the inductance  $L$ , and the frequency of the current  $\omega$  according to  $Z = \sqrt{R^2 + \left(\omega L - \frac{1}{\omega C}\right)^2}$ . Solve this equation for the inductance  $L$ .
47. The formula used to find the orbital period for circular Keplerian orbits is  $P = \frac{2\pi}{\sqrt{\frac{u}{a^3}}}$ . Solve this equation for  $a$ .