

Chapter 5

Review Exercises

1–2 Use $(O_4 + U_4)/2$ to estimate the area under the graph of the function and above the x -axis on the given interval.

1. $f(x) = \frac{x^2}{2}$ on $[0, 2]$

2. $f(x) = \sin x$ on $\left[0, \frac{\pi}{2}\right]$

3–6 Write the given sum using sigma notation.

3. $\frac{1}{2} - \frac{1}{9} + \frac{1}{28} - \frac{1}{65} + \cdots - \frac{1}{1,000,001}$

4. $a_1 + a_5 + a_9 + a_{13} + \cdots + a_{97}$

5. $f\left(\frac{2}{n^2}\right) + f\left(\frac{4}{n^2}\right) + f\left(\frac{6}{n^2}\right) + \cdots + f\left(\frac{100}{n^2}\right)$

6. $g(t_{-2}^*)\Delta t + g(t_{-1}^*)\Delta t + g(t_0^*)\Delta t + \cdots + g(t_{2n}^*)\Delta t$

7–8 Assuming that $\sum_{i=0}^n a_i = 50$ and $\sum_{i=0}^n b_i = 80$, find the sum.

7. $\sum_{i=0}^n (a_i + 2b_i + 2)$

8. $\sum_{i=0}^n \left(\frac{a_i}{5} - \frac{b_i}{4}\right)$

9–10 Use summation formulas to find the value of the sum.

9. $\sum_{i=1}^{10} (3i^3 - 1)$

10. $\sum_{j=1}^n \frac{(2j+1)(j-2)}{2}$

11. Find the value of the sum $\sum_{i=1}^n \left[\frac{1}{i^2} - \frac{1}{(i+1)^2} \right]$.

(Hint: Write out the first few terms as well as the last few terms.)

12–13 Evaluate the geometric sum using the formula proven in Exercise 53 of Section 5.1.

12. $\sum_{i=0}^{10} \frac{3}{2^i}$

13. $\sum_{j=0}^6 (-1)^j (0.3)^j$

14–15 Evaluate the given double sum.

14. $\sum_{i=3}^{11} \sum_{j=1}^4 (2i - j)$

15. $\sum_{i=1}^n \sum_{j=1}^m i^2 j$

16–18 Find the area under the graph of $f(x)$ and above the x -axis on the given interval, by taking the limit of the associated Riemann sums.

16. $f(x) = x^2 + 1$ on $[0, 2]$

17. $f(x) = x^3$ on $[0, 1]$

18. $f(x) = \sqrt{x}$ on $[0, 4]$

(Hint: Choose $x_i^* = \frac{4(i-1)^2}{n^2}$.)

19. Identify the region whose area is the limit given by

$\lim_{n \rightarrow \infty} \frac{2}{n} \sum_{i=1}^n \left[\frac{4i}{n} - \left(\frac{2i}{n} \right)^2 \right]$. Then use summation formulas to evaluate the limit.

20–21 Prove that the given function is not integrable on the interval $[0, 1]$.

20. $f(x) = \frac{1}{x^2}$

21. $g(x) = \begin{cases} 1 & \text{if } x \text{ is rational} \\ -1 & \text{if } x \text{ is irrational} \end{cases}$

22–25 Suppose that f is an even function, g is odd, both are integrable on $[-a, a]$, and we know that $\int_0^a f(x) dx = 2$, while $\int_0^a g(x) dx = 0.5$ ($a > 0$). If possible, find the integral.

22. $\int_{-a}^a [5f(x) + 4g(x)] dx$

23. $\int_{-a}^a [f(x)]^2 g(x) dx$

24. $\int_{-a}^a f(x) [g(x)]^2 dx$

25. $\int_{-a}^0 [f(x) + g(x)] dx$

26–27 Find the average value of $f(x)$ over the given interval and identify all points in the domain where $f(x)$ assumes its average value.

26. $f(x) = 4x - x^2$ on $[0, 4]$

27. $f(x) = |x - 2| - 1$ on $[1, 5]$

28. Use the Fundamental Theorem of Calculus to evaluate the limit

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \left(\sqrt{\frac{i}{n}} - \frac{i}{n} \right)$$

by recognizing it as a Riemann sum of a function over an interval.

- 29–30 Use Part I of the Fundamental Theorem of Calculus to find the derivative of the given function.

29. $F(x) = \int_0^x \sqrt{1+t^2} dt$ 30. $G(x) = \int_0^{x^2} e^{t^2} dt$

- 31–38 Use Part II of the Fundamental Theorem of Calculus to evaluate the definite integral.

31. $\int_1^2 (2x^4 + 3x^2 - 2) dx$ 32. $\int_0^2 (3x+2)(5-x) dx$

33. $\int_1^4 \left(\frac{1}{t} - \frac{2}{t^2} + 1 \right) dt$ 34. $\int_1^9 \frac{x^2 - 2\sqrt{x} + 2}{x} dx$

35. $\int_0^1 \frac{2}{\sqrt{1-x^2}} dx$

36. $\int_{\pi/4}^{3\pi/4} (2 \csc^2 x - \cos x) dx$

37. $\int_2^3 \frac{x+2}{x-1} dx$ 38. $\int_0^1 \frac{2x^2-1}{x^2+1} dx$

- 39–40 Find the area of the region between the graph of the given function and the x -axis on the indicated interval.

39. $y = \frac{1}{2x^2}$ on $[1, 10]$

40. $y = 2\sqrt{x} - x^2$ on $[0, 2]$

41. Find a formula for $f(x)$ if $\int_0^{x^3} f(t) dt = \sin(x^3)$.

42. The velocity function of a particle moving along the x -axis is $v(t) = 3t - t^2$ units per second. If it started at the origin, find **a.** the position of the particle at $t = 5$ seconds and **b.** the total distance traveled by the particle in the time interval $[0, 5]$.

- 43–54 Use an appropriate substitution (when necessary) to evaluate the indefinite integral.

43. $\int \frac{-2}{\sqrt{1-x^2}} dx$ 44. $\int \frac{-2x}{\sqrt{1-x^2}} dx$

45. $\int \sec x (\sec x + \tan x) dx$

46. $\int \sec^2 x \tan x dx$

47. $\int 6x^2 (2x^3 - 7)^9 dx$ 48. $\int x^4 \sqrt{x^5 - 3} dx$

49. $\int \frac{4x}{(x^2+1)^2} dx$

50. $\int \frac{e^{\arcsin x}}{\sqrt{1-x^2}} dx$

51. $\int \frac{\sec^2 \sqrt{x}}{\sqrt{x}} dx$

52. $\int \frac{1}{x \ln(x^2)} dx$

53. $\int \frac{e^{2x}}{e^x+1} dx$

54. $\int \frac{1}{x^2} \sin\left(\frac{x+1}{x}\right) dx$

- 55–56 Find $y(x)$ that satisfies the given conditions.

55. $\frac{dy}{dx} = \frac{1}{\sqrt{x}(\sqrt{x}-1)^2}$; $y(9) = 1$

56. $y''(x) = 1 - \sin x$; $y'(0) = 2$; $y(0) = 0$

57. A particle is moving along the x -axis in the positive direction with a velocity function of $v(t) = \frac{t}{t^2+1}$ units per second. If it started at the point $(1, 0)$, what is the particle's position at $t = 4$ seconds?

- 58–63 Evaluate the definite integral.

58. $\int_0^1 3x^5 (x^6 - 1)^{12} dx$

59. $\int_0^4 (x+1)\sqrt{x^2+2x} dx$

60. $\int_1^2 \frac{x^3}{x^4+1} dx$

61. $\int_{-1/2}^0 \frac{2^t}{\sqrt{1-4^t}} dt$

62. $\int_e^{e^2} \frac{1}{x(\ln x)^2} dx$

63. $\int_1^4 \frac{dx}{\sqrt{x}(\sqrt{x}+1)^2}$

- 64–69 Find the area of the region bounded by the graphs of the given equations. (If convenient or necessary, integrate with respect to y rather than x .)

64. $y = x^3 - 4x$, $3y = 15x$, $x \geq 0$

65. $y = 1 - x^2$, $y = 1 - x^6$, $x \geq 0$

66. $y = 2\sqrt{x}$, $y = 4 - 2x$, $y = 0$

67. $y = \ln x$, $(e-1)y = x - 1$

68. $y = \frac{1}{1+x^2}$, $2y = 1$

69. $y = \sin x$, $y = \sin x \cos x$, $0 \leq x \leq \pi$

70. Consider the region bounded by the graph of $y = 1/x$ and the x -axis over the interval $[1, a]$ ($a > 1$). Find the vertical line $x = c$ that bisects the region in two subregions of equal area.

71. Consider the function $f(x) = 1/x^2$ defined on some interval $[a, b]$. Partition $[a, b]$ and in each subinterval $[x_{i-1}, x_i]$ choose the sample point $x_i^* = \sqrt{x_{i-1}x_i}$ (the geometric mean of the endpoints). Show that

$$\frac{1}{(x_i^*)^2} \Delta x_i = \frac{1}{x_{i-1}} - \frac{1}{x_i}$$

and use this observation to prove the following formula.

$$\int_a^b \frac{1}{x^2} dx = \frac{1}{a} - \frac{1}{b}$$

72. Prove that if the conditions of Part I of the Fundamental Theorem of Calculus are satisfied and $F(x) = \int_{g(x)}^{h(x)} f(t) dt$, where $g(x)$ and $h(x)$ are differentiable, then $F'(x) = f(h(x))h'(x) - f(g(x))g'(x)$. (Hint: See Example 3 of Section 5.3.)
73. Prove that if f is a linear function, then its definite integral on an interval $[a, b]$ is the average of its left and right Riemann sums, that is,

$$\int_a^b f(x) dx = \frac{L_n + R_n}{2}.$$

What is your expectation regarding the integral and the average above if f is concave up? Concave down?

Concept Check

- 74–81 Determine whether the given statement is true or false. In case of a false statement, explain or provide a counterexample.
74. If $n_1 < n_2$, then the Riemann sum R_{n_2} is always a better approximation of the integral than R_{n_1} .

75. If f is piecewise continuous on a closed interval, then the limit of its Riemann sums always exists.
76. When applying the Fundamental Theorem of Calculus, we must choose the antiderivative with $C = 0$.
77. $\int \frac{1}{e^x} dx = \ln(e^x) + C = x + C$
78. The definite integral of the velocity function of a moving object on $[t_1, t_2]$ is equal to the total distance traveled by the object from time $t = t_1$ to $t = t_2$.
79. $\int_a^b f(x) dx > 0$ if and only if $f(x) > 0$ on $[a, b]$.
80. $\int \sec x dx = \sec x \tan x + C$
81. $\int_{-1}^1 \frac{1}{x^3} dx = \frac{-1}{2x^2} \Big|_{-1}^1 = -\frac{1}{2} - \left(-\frac{1}{2}\right) = 0$

Chapter 5 Technology Exercises

82. Use the summation feature of a graphing utility to verify your answers for Exercises 9–15.
83. Write a program for a graphing calculator or computer algebra system that calculates the n^{th} Riemann sum for a given function on a given interval, using subintervals of equal width and sample points of your choice. Use your program to verify your answers for Exercises 16–18.
84. Use a graphing utility to evaluate the limit of Exercise 28. What do you find? (Answer will vary depending on the capabilities of the particular software used.)