

Thus $x = 0$ and $x = \frac{4}{3}$ will make $f''(x)$ equal to zero. Both values give inflection points for $f(x)$.

We can use $f'(x) = 0$ to locate the maximum and minimum points for $f(x)$.

$$\begin{aligned} f'(x) &= 0 \\ 8x^2 - 4x^3 &= 0 \\ 4x^2(2-x) &= 0 \\ x &= 0, 2 \end{aligned}$$


In Figure 8, we observe $x = 0$ corresponds to an inflection point (neither a maximum nor a minimum) and $x = 2$ corresponds to the maximum point $\left(2, \frac{31}{3}\right)$.

In this problem, the algebraic solutions of $f'(x) = 0$ and $f''(x) = 0$ are of a familiar type. When these equations are too difficult for an algebraic solution, a graphing utility is invaluable.

12.1 EXERCISES

💡 PRACTICE

- At each point marked on the graph of f , determine if f' is positive, negative, or zero. Determine if f'' is positive, negative or zero.







Draw a graph that satisfies the given conditions in Exercises 2–5.

- $f(5) = 9$, $f'(5) = 2$, $f''(5) = -2$
- $f(-5) = -9$, $f'(-5) = 2$, $f''(-5) = 2$
- $f(5) = -9$, $f'(5) = 0$, $f''(5) = 3$
- $f(0) = 12$, $f'(0) = 0$, $f''(0) = -3$



For Exercises 6–13, find $f''(x)$. Then evaluate $f''(0)$, $f''(1)$, and $f''(4)$, if they exist.



- $f(x) = x^3 + x^2 + 3$
- $f(x) = x^3 - x^2 + 7$
- $f(x) = x^2 - 5\sqrt{x} + 1$
- $f(x) = x^2 + 2\sqrt{x} - 3$
- $f(x) = \sqrt{x-4}$
- $f(x) = \sqrt{2x+1}$
- $f(x) = \frac{x}{x+5}$
- $f(x) = \frac{x-2}{x+4}$

In Exercises 14–17, sketch a possible graph for $f'(x)$ on the same coordinate axes as $f(x)$. Then locate all inflection points on the graph of $f(x)$.



For each of the graphs in Exercises 18–21, list the interval(s) **a.** on which f is concave upward and **b.** on which f is concave downward; then **c.** locate all points of inflection.





In Exercises 22–33, determine the intervals on which each function is **a.** concave upward and **b.** concave downward; then **c.** locate all points of inflection. Use the information gathered to sketch the function. Confirm the details with a graphing calculator.

22. $f(x) = 2x^2 + 5x - 9$

23. $f(x) = 5x^2 + 8x - 1$

24. $f(x) = x^3 - 3x^2 + 7$

25. $f(x) = x^3 + 6x^2 - 10$

26. $f(x) = x^3 + 11x - 4$

27. $f(x) = 5x^3 + 7x + 2$

28. $f(x) = \frac{1}{3}x^3 - 2x^2 + x - 3$

29. $f(x) = \frac{1}{3}x^3 + 3x^2 + 2x - 5$

30. $f(x) = \sqrt[3]{2x + 3}$

31. $f(x) = \sqrt[3]{5x - 3}$

32. $f(x) = \frac{x}{x^2 - 4}$

33. $f(x) = \frac{4x}{x^2 - 5}$

WRITING & THINKING

In Exercises 34–37, give an example of a polynomial function that satisfies the conditions.


34. $F(5) = 15$; $F'(x)$ is nonzero, but $F''(x) = 0$ for all x .

35. $G(0) = 0$, $G'(0) = 0$, and $G''(0) = 0$; $G(x)$ is concave upward everywhere and has no inflection points.

36. $H(4) = 0$; $H'(x)$ is positive for $x > 4$ and negative for $x < 4$. $H(x)$ has no inflection points.

37. $J(4) = 0$; $J'(4)$ is zero but $J'(x)$ is positive if $x \neq 4$; $J''(4) = 0$.

APPLICATIONS



38. **Filtrate:** In a chemistry lab a filtrate drips slowly but continuously at a constant rate into a glass container shaped like the one shown. The container eventually fills to the base of the neck. Let t denote the passage of time and h be the height of the liquid.

- Describe at what points on the bottle $\frac{dh}{dt}$ will be a maximum and a minimum.
- Sketch a graph of $\frac{dh}{dt}$. Are there any inflection points on a graph of $y = h(t)$?
- Add a sketch of $y = h(t)$ on the same coordinate axes as in part **b**.