

## 10.R.4 Exercises

### Concept Check

**True/False.** Determine whether each statement is true or false. If a statement is false, explain how it can be changed so the statement will be true. (**Note:** There may be more than one acceptable change.)

---

1. When a decimal number is changed to a fraction, the denominator will be the power of 10 that names the rightmost digit of the decimal number.
2. When a decimal number is changed to a fraction, the numerator can be determined by using the whole number that is formed by all the digits of the decimal number.
3. Fractions can always be converted to decimal form without losing accuracy.
4. In decimal form,  $\frac{1}{3}$  is repeating and nonterminating.

### Practice

Change each decimal number to a fraction or mixed number in lowest terms.

---

5. 0.18

6. 2.75

Change each fraction to a decimal number rounded to the nearest hundredth.

---

7.  $\frac{20}{3}$

8.  $\frac{40}{9}$

Simplify the expression by first writing all of the numbers in decimal form. Round to the nearest hundredth, if necessary.

---

9.  $\frac{1}{4} + 0.25 + \frac{1}{5}$

10. Arrange  $0.76$ ,  $\frac{3}{4}$ ,  $\frac{7}{10}$  in order from smallest to largest.

## Applications

Solve.

---

11. A rectangle measures 6.4 inches in length, and has a width that measures  $\frac{2}{5}$  of the length. Find the perimeter of the rectangle.
12. A loaf of bread weighs 21.6 ounces. Mauricio cut off a third of the loaf to save for later and then cut the remaining portion into 16 equal slices. What was the weight of each slice of the 16 slices he cut?

## Writing & Thinking

13. Describe the process used to change a terminating decimal number to a fraction.

14. List 2 different ways to solve this problem:  $\frac{1}{2} + 3.67 - \frac{1}{8}$ . State which method you prefer and why.